

Blended Feeding Fats

Typical Analysis

Moisture	0.75%
Total Fatty Acids	90.00%
Free Fatty Acids	15.00%
Impurities (I)	0.10%
Unsaponifiables (U)	0.65%
Total MIU	1.50%
FAC Color	39
Peroxide Value	<5%
Viscosity	35.0 CST @ 120°F
Weight	7.44 lb/gal
Unsaturated to Saturated Ration	2.68:1
Iodine Value (IV)	79.00%
Titre	32.50°C
Metabolizable Energy	3,800 Kcal/lb = 8,377 Kcal/kg

Minerals & Vitamins

C8 Caprylic Acid	--	C17 Margaric Acid	0.15%
C10 Capric Acid	--	C18 Stearic Acid	10.78%
C12 Lauric Acid	--	C18:1 Oleic Acid	51.03%
C12:1 Lauroleic Acid	--	C18:2 Linoleic Acid	17.86%
C13 Tridecanoic Acid	--	C 18:3 Linolenic Acid	1.25%
C14:1 Myristoleic Acid	0.15%	C20:1 Eicosenoic	0.45%
C15 Pentadecanoic	0.01%	C22 Behenic Acid	0.08%
C16 Palmitic Acid	15.18%	C22:1 Erucic	0.08%
C16:1 Palmitoleic Acid	1.36%	C24 Lignoceric Acid	0.01%
C16:2 Hexadecadienoic Acid	0.36%	C24:1 Tetracosenoic Acid	0.26%

Analysis are updated periodically and may vary slightly from previous versions.

Approved plant of:

Animal Protein Producers Institute (APPI), Voluntary Salmonella Reduction Program
Food & Drug Administration (FDA)
Animal & Plant Health Inspection Service (APHIS)
Hazard Analysis & Critical Control Point (HACCP)

